

Use of exogenous gonadotropins in anovulatory women: a technical bulletin

The Practice Committee of the American Society for Reproductive Medicine

American Society for Reproductive Medicine, Birmingham, Alabama

This Technical Bulletin reviews the indications for gonadotropin treatment in anovulatory women; outlines the recommended pretreatment evaluation, treatment regimens, and monitoring; describes the alternatives and adjuncts to treatment with gonadotropins; and summarizes the complications of gonadotropin treatment. (*Fertil Steril*® 2008;90:S7–12. ©2008 by American Society for Reproductive Medicine.)

Exogenous gonadotropins can be used for ovulation induction (OI) in women who are anovulatory and infertile, and they are indicated when OI cannot be achieved with less complex methods (1). The purpose of this document is to review the indications for gonadotropin treatment in anovulatory women; to outline the recommended pretreatment evaluation, treatment regimens, and monitoring; to describe briefly the alternatives and adjuncts to treatment with gonadotropins; and to summarize the complications of gonadotropin treatment. The use of gonadotropins to purposely induce superovulation in concert with intrauterine insemination or assisted reproductive technology (ART) is not considered or discussed in this document, because the principles, mechanisms, and goals of such treatments are distinctly different from those of OI.

INDICATIONS

Gonadotropins are used to treat infertility caused by anovulation, usually after other less complicated and costly methods have failed. Two groups of anovulatory disorders may require gonadotropin therapy: 1) hypogonadotropic amenorrhea (HA, also known as hypogonadotropic hypogonadism, hypothalamic amenorrhea, or World Health Organization [WHO] type I amenorrhea); and 2) polycystic ovary syndrome (PCOS, also known as hyperandrogenic amenorrhea or WHO type II amenorrhea) (2, 3). Premature ovarian “failure” (POF) (also known as hypergonadotropic hypogonadism or WHO type III amenorrhea) is not responsive to exogenous gonadotropins and must be excluded (2, 3).

Hypogonadotropic amenorrhea (HA)

Endogenous circulating FSH and LH concentrations are normal or low in women with HA. Although pituitary disorders, such as panhypopituitarism, can cause HA, the disorder usually results from very low or absent hypothalamic GnRH se-

cretion (2, 3). Such disorders of endogenous GnRH secretion can be congenital or acquired and may relate to severe weight loss (anorexia nervosa), excessive chronic physical exercise, severe emotional stress, or medications, or may be idiopathic (2, 3). Hyperprolactinemia resulting from a pituitary adenoma also can result in secondary or acquired HA. The possibility should be investigated specifically and treated when identified with a dopamine agonist before OI, because ovulatory function will be restored in most women after PRL levels fall into the normal range (4). Women with HA seldom respond to clomiphene citrate (CC); alternative methods for OI include pulsatile GnRH (5) and exogenous gonadotropins. However, gonadotropin treatment should include both FSH and LH activity to effectively stimulate both steroidogenesis and folliculogenesis (6).

Polycystic ovary syndrome (PCOS)

Women with PCOS generally have normal or low serum FSH and often mildly increased LH concentrations. Although most will respond to treatment with CC, many do not, particularly if obese (7). Exogenous gonadotropin treatment often is required in CC-resistant women with PCOS, but it is also associated with significantly increased risk for ovarian hyperstimulation syndrome (OHSS) and multiple pregnancy (7, 8). Accordingly, low-dose gonadotropin treatment regimens are strongly advised (7).

PRETREATMENT EVALUATION

Gonadotropin therapy is costly and has significant risks, and should therefore be used only by clinicians having the requisite training and experience. Before treatment begins, other potential coexisting causes of infertility, such as uterine cavity abnormalities (myomas, adhesions), tubal obstruction, advanced endometriosis (ovarian endometrioma), pelvic adhesions, or poor semen quality, should be excluded. Pretreatment evaluation generally should exclude abnormalities of thyroid function and hyperprolactinemia and should include hysterosalpingography, transvaginal ultrasonography, and semen analysis. Women with POF should not be considered as candidates for OI with exogenous gonadotropins (2).

Technical Bulletin

Received July 28, 2008; accepted August 5, 2008.

No reprints will be available.

Correspondence: Practice Committee, American Society for Reproductive Medicine, 1209 Montgomery Highway, Birmingham, AL 35216.

GONADOTROPIN PREPARATIONS

Gonadotropin products for human use derive from urinary extracts or recombinant technology and all have similar effectiveness and safety (9). Detailed information on the composition and pharmacology of gonadotropin preparations is provided in a separate document entitled "Gonadotropin Preparations: Past, Present, and Future Perspectives" (10).

GONADOTROPIN REGIMENS FOR OI

Exogenous FSH stimulates proliferation of granulosa cells and follicular growth. Luteinizing hormone stimulates the production of androgen in thecal cells that subsequently is converted to estrogen by granulosa cells via aromatization (1). The goal of treatment is to promote the growth and development of a single mature follicle.

In women with PCOS, OI usually begins after a menses induced by a brief interval of treatment with an exogenous progestogen. In women with HA, OI may begin at any convenient time. Baseline ultrasonography is prudent to exclude ovarian cysts that might be confused with new follicular growth. Among women not previously treated with exogenous gonadotropins, treatment generally should begin at a relatively low dose (e.g., 37.5–75 IU/day). In subsequent cycles, treatment generally begins at the threshold of response previously determined. Although 7–12 total days of treatment is typical, longer durations of treatment may be required. Once a mature follicle has developed, generally measuring 16–18 mm in mean diameter, exogenous hCG or a GnRH agonist is administered to stimulate ovum release, as described in a subsequent section (1, 7).

Monitoring of OI

The safety and efficacy of gonadotropin treatment depend on careful monitoring with serial transvaginal ultrasonography and serum E₂ measurements (1, 7). Ultrasonography provides a structural measure of follicular development and generally should be performed after the first 4–5 days of treatment and at intervals of 1–3 days thereafter according to response (1, 7). Endometrial thickness and appearance provide an indirect measure of endometrial development and maturation and have some prognostic value for implantation (1, 11). Rapid serum E₂ determinations (results available the same day and, ideally, 7 days/week) provide a functional measure of follicular maturation. Together with ultrasonography, serum E₂ levels provide an accurate gauge of response to treatment and guide the management of the OI cycle (1, 7).

Gonadotropin Regimens in HA

In women with HA, optimal clinical results are achieved by the combined administration of FSH and LH (12, 13), accomplished by administration of hMG (14) or a combination of FSH and either recombinant LH (15, 16) or low-dose hCG (17). In addition to stimulating the production of intrafollicular androgens which provide the substrate for estrogen pro-

duction that enhances oocyte development, LH activity promotes development of larger follicles (16–19). There are no established gonadotropin regimens for OI of HA patients and no prospective studies that assess treatment outcomes with different gonadotropin dosages.

Gonadotropin Regimens in PCOS

In women with PCOS, only FSH activity is required, because endogenous LH levels are adequate, although added LH does not appear to be harmful (1, 7). There is no significant advantage to using any specific gonadotropin preparation. A meta-analysis concluded that the outcomes of treatment achieved with hMG and with FSH alone were similar (20). Others have observed that treatment with rFSH or urinary FSH yields similar results (21).

The risk of OHSS and multiple pregnancy is greater among women with PCOS than among those with HA, primarily because gonadotropin treatment generally stimulates development of larger follicular cohorts in women with PCOS (22). Consequently, exogenous gonadotropins must be administered judiciously (23–29). The recommended approach in the first dose-finding cycle is to begin with a low dose of gonadotropin, typically 37.5–75 IU/day, increasing after 7 days or more if no follicle >10 mm has yet emerged, in small increments, at intervals, until evidence of progressive follicular development is observed. The maximum required daily dose of FSH/hMG seldom exceeds 225 IU/day (23).

Inducing Oocyte Release

The final stages of follicle/oocyte maturation and ovum release can be induced by administration of hCG extracted from urine, recombinant hCG, recombinant LH, or a GnRH agonist (1, 7, 30–33). Although there are no evidence-based specific guidelines for optimal timing, the ovulatory stimulus generally should be administered when at least one and, ideally, no more than two follicles greater than 16–18 mm in mean diameter are observed, so as to limit the risk for multiple pregnancy. Estradiol concentrations generally should range between approximately 150 and 300 pg/mL per dominant follicle.

Human chorionic gonadotropin is administered in a single injection of 5,000–10,000 IU IM or SC (1, 7); recombinant hCG is administered at a dose of 250 µg SC, which corresponds to approximately 6,000–7,000 IU hCG (30). Recombinant LH also is effective and potentially safer than hCG. However, the current formulation of recombinant LH (75 IU/vial) precludes its use for that purpose, because 25,000–30,000 IU recombinant LH is required to yield results similar to those achieved with 5,000 IU hCG (31). Given these considerations, hCG is the least expensive and most widely used preparation (1).

A GnRH agonist also can be used to induce oocyte release in OI, provided that an agonist has not been administered earlier during the course of gonadotropin treatment to prevent

a spontaneous LH surge. The GnRH agonists are not effective in HA. For purposes of inducing ovum release, the recommended dose for leuprolide is 500 μg (32) and for triptorelin is 200 μg (33); both are administered in a single SC injection. When a GnRH agonist is used to induce ovum release, progesterone supplementation during the luteal phase is required, because agonist treatment may adversely affect endogenous luteal function.

Support of Corpus Luteum Function

Routine treatment with supplemental exogenous progesterone or additional small doses of hCG (1,500–2,500 IU every 3–4 days) during the luteal phase of OI cycles is advocated by many. Luteal support after OI is most clearly indicated and recommended in women with HA whose endogenous gonadotropin secretion may be inadequate to support normal luteal function. Its necessity and effectiveness otherwise are unproven and controversial (1).

TIMING OF INTERCOURSE

Ovulation can be expected to occur between 24 and 48 h after injection of hCG or a GnRH agonist. Consequently, intercourse within that interval can help to maximize the likelihood of conception. When specifically indicated, intrauterine insemination generally should be planned for approximately 24–36 h after the injection (1).

RESULTS ACHIEVED WITH GONADOTROPIN TREATMENT FOR OI

In a 2003 systematic review of 13 studies published between 1991 and 2000 including 1,269 cycles among 881 anovulatory infertile women (2.7 cycles per patient) who received gonadotropin treatment for OI, there were 366 pregnancies; the pregnancy rate was 15% per cycle and 41% per patient (34). Women who were obese or insulin resistant required increased total amounts of gonadotropins per cycle. Insulin resistance, but not obesity, was associated with a lower pregnancy rate (odds ratio [OR] 0.24, 95% confidence interval [CI] 0.08–1.71). Another study, not included in the 2003 systematic review, reported 18 pregnancies in 28 women with HA across 72 cycles of treatment (25% per cycle), but only 16 pregnancies in 61 hyperandrogenic women with PCOS across 205 cycles of treatment (8% per cycle) (35). The type of gonadotropin did not appear to have significant influence on pregnancy rate in four trials involving 396 patients treated with FSH (12%) or hMG (10%) (OR 0.82, 95% CI 0.44–1.53) (36).

ALTERNATIVES AND ADJUNCTS TO GONADOTROPIN TREATMENT FOR OI

Clomiphene citrate (CC) is traditional first-line treatment for OI. Clomiphene citrate competes with endogenous estrogen for the estrogen receptor and, by causing a central misinterpretation of circulating estrogen levels, induces a transient but significant increase in the secretion of pituitary FSH. In

women with HA in whom the hypothalamic-pituitary-ovarian feedback axis is severely dysfunctional, CC is ineffective. Clomiphene citrate may also be used in concert with exogenous gonadotropins in a sequential regimen that begins with CC, adding gonadotropins on the last day of CC therapy or the day after. Limited evidence suggests that sequential CC/gonadotropin therapy may help to lower the total dose, duration, and cost of gonadotropin treatment required to achieve OI (37, 38).

Metformin enhances insulin actions and has been advocated for OI in women with PCOS, in whom the prevalence of insulin resistance is relatively high. Although some anovulatory women will ovulate in response to metformin treatment alone (1,500 mg/day), CC has proven to be more effective in a large randomized clinical trial (39). Metformin treatment may be effective in nonobese women with PCOS (40). Combined treatment with metformin and FSH also has been advocated. Although combined treatment may help to reduce the total dose of gonadotropins required to achieve OI (41), pregnancy rates are not increased (42). The clinical value of metformin in CC-refractory anovulatory women who require gonadotropin therapy has not been investigated thoroughly.

Ovarian drilling has been used to decrease ovarian androgen production in the ovaries of anovulatory infertile women with PCOS and can successfully induce ovulation in many such women who may prove resistant to treatment with CC (43, 44). However, the benefits that ovarian drilling may offer are usually transient and the procedure has risks, such as pelvic adhesions and potential adverse effects on ovarian reserve (43, 44). In highly selected individuals, ovarian drilling may be an alternative to gonadotropin treatment that merits consideration. However, even after ovarian drilling, OI still may be required. With the ready availability of ART in most centers, ovarian drilling has become uncommon, although the procedure can achieve good results and is associated with a lower risk for multiple pregnancy (43, 44).

Gonadotropin-releasing hormone analogs (agonists and antagonists) are most commonly used in conjunction with gonadotropin treatment in ART stimulation cycles, primarily to prevent premature luteinization and ovulation (1). Although seldom used in OI, they may have some utility for individuals in whom those phenomena are observed during gonadotropin treatment. When used in OI cycles, treatment with supplemental progesterone during the luteal phase generally is recommended because endogenous pituitary LH secretion remains suppressed for some time after treatment with a GnRH agonist (45) or antagonist (46, 47).

Aromatase inhibitors block conversion of androgens to estrogens. When normally operating, the feedback relationship between circulating estrogen levels and gonadotropin secretion results in an increase in pituitary FSH secretion (48). No trials have evaluated aromatase inhibitors as an alternative to gonadotropin treatment in CC-resistant women. When used in sequence or in conjunction with exogenous

gonadotropins, aromatase inhibitors may reduce the amount and duration of gonadotropins required to achieve OI, but E₂ levels are more difficult to interpret confidently, and the evidence to support their use is meager (49). Aromatase inhibitors are not approved for use in OI. Experience with the off-label use of aromatase inhibitors for OI is limited (50), their safety has been questioned (51, 52), and studies of their efficacy and clinical utility are ongoing.

COMPLICATIONS IN OI

Multifetal gestation is the most frequent complication of OI. The goal of OI is ovulation of a single mature oocyte, and although that objective can sometimes be difficult to achieve (53), the criteria for cycle cancellation should nonetheless be stringent. To minimize the risk of multifollicular ovulation and multiple pregnancy, cycle cancellation generally should be seriously considered when three or more mature follicles (>16–17 mm) or large numbers of intermediate-sized follicles (10–15 mm) are observed or when the serum E₂ concentration exceeds 1,000–1,500 pg/mL (1). However, the increased risk for multiple pregnancy associated with gonadotropin treatment cannot be eliminated entirely, and even the low-dose gonadotropin regimens advocated here are associated with five- to 10-fold increased risk (54). Although multifetal pregnancy reduction can be offered when treatment results in a high-order multiple pregnancy (triplets or greater), the risk is easier to avoid than to manage (55).

Ovarian hyperstimulation syndrome (OHSS) can occur after OI in anovulatory women, and the risk cannot be eliminated completely. To minimize the risk for developing OHSS, a lower dose of hCG (5,000 IU) or a GnRH agonist is recommended for inducing ovum release when E₂ levels have risen rapidly or are markedly elevated (>2,500 pg/mL) and/or a large number of intermediate-sized follicles (10–14 mm) are observed (56).

Earlier concerns that OI might be associated with an increased risk for cancer of the ovary and breast (57) have not been corroborated by subsequent studies (58, 59). Although the risk for ovarian cancer may be higher for infertile women than for fertile women, there is no compelling evidence to indicate that such risk is increased by OI.

SUMMARY AND RECOMMENDATIONS

- Gonadotropin treatment for OI is indicated for anovulatory infertile women with HA and for those with PCOS who fail to respond to less complicated OI treatment regimens.
- Before treatment with gonadotropins, evaluation generally should exclude abnormalities of thyroid function and hyperprolactinemia, and should include hysterosalpingography, transvaginal ultrasonography, and evaluation of the male partner by semen analysis.
- Although sometimes difficult to achieve, the goal of gonadotropin treatment is to promote the growth and development of a single mature follicle.

- Among women not previously treated with exogenous gonadotropins, treatment generally should begin at a relatively low dose (e.g., 37.5–75 IU/day). In subsequent cycles, treatment generally begins at the threshold of response previously determined.
- Ovulation of a mature ovarian follicle may be triggered with either purified (5,000–10,000 IU IM or SC) or recombinant (250 µg SC) hCG. A GnRH agonist (leuprolide 500 µg SC; triptorelin 200 µg SC) also may be used to trigger ovulation, except in women with HA and those who have received GnRH agonist treatment earlier during the course of gonadotropin treatment.
- Luteal support after OI is most clearly indicated and recommended in women with HA and those who receive treatment with a GnRH agonist, because, in these cases, endogenous gonadotropin secretion may be inadequate to support normal luteal function.
- The increased risk of multiple pregnancy associated with gonadotropin treatment can be limited by judicious treatment and careful monitoring, but it cannot be eliminated entirely.
- Despite careful monitoring, OHSS can occur after OI in anovulatory women.

Acknowledgments: This report was developed under the direction of the Practice Committee of the American Society for Reproductive Medicine as a service to its members and other practicing clinicians. Although this document reflects appropriate management of a problem encountered in the practice of reproductive medicine, it is not intended to be the only approved standard of practice or to dictate an exclusive course of treatment. Other plans of management may be appropriate, taking into account the needs of the individual patient, available resources, and institutional or clinical practice limitations. The Practice Committee of the American Society for Reproductive Medicine and the Board of Directors of the American Society for Reproductive Medicine have approved this report.

REFERENCES

1. Huime JA, Lambalk CB, van Loenen AC, Schats R, Hompes PG, Fauser BC, et al. Contemporary pharmacological manipulation in assisted reproduction. *Drugs* 2004;64:297–322.
2. Practice Committee of the American Society for Reproductive Medicine. Current evaluation of amenorrhea. *Fertil Steril* 2006;86(Suppl 5): S148–55.
3. Reindollar RH, Novak M, Tho SP, McDonough PG. Adult-onset amenorrhea: a study of 262 patients. *Am J Obstet Gynecol* 1986;155:531–43.
4. Biller BM, Luciano A, Crosignani PG, Molitch M, Olive D, Rebar R, et al. Guidelines for the diagnosis and treatment of hyperprolactinemia. *J Reprod Med* 1999;44(Suppl 12):1075–84.
5. Filicori M. Clinical review 81: gonadotropin-releasing hormone analogs in ovulation induction: current status and perspectives. *J Clin Endocrinol Metab* 1996;81:2413–6.
6. Burgues S, Spanish Collaborative Group on Female Hypogonadotropic Hypogonadism. The effectiveness and safety of recombinant human LH to support follicular development induced by recombinant human FSH in WHO group I anovulation: evidence from a multicentre study in Spain. *Hum Reprod* 2001;16:2525–32.
7. Cristello F, Cela V, Artini PG, Genazzani AR. Therapeutic strategies for ovulation induction in infertile women with polycystic ovary syndrome. *Gynecol Endocrinol* 2005;21:340–52.
8. Ratts VS, Pauls RN, Pinto AB, Kraja A, Williams DB, Odem RR. Risk of multiple gestation after ovulation induction in polycystic ovary syndrome. *J Reprod Med* 2007;52:896–900.

9. Bayram N, van Wely M, van Der Veen F. Recombinant FSH versus urinary gonadotropins or recombinant FSH for ovulation induction in subfertility associated with polycystic ovary syndrome. *Cochrane Database Syst Rev* 2001;(2):CD002121.
10. The Practice Committee of the American Society for Reproductive Medicine. Gonadotropin preparations: past, present, and future perspectives. *Fertil Steril* 2008;90 (Suppl 3):S13–20.
11. Baruffi RL, Contart P, Mauri AL, Petersen C, Felipe V, Garbellini E, et al. A uterine ultrasonographic scoring system as a method for the prognosis of embryo implantation. *J Assist Reprod Genet* 2002;19:99–102.
12. Filicori M, Cognigni GE, Pocognoli P, Tabarelli C, Ferlini F, Perri T, et al. Comparison of controlled ovarian stimulation with human menopausal gonadotropin or recombinant follicle-stimulating hormone. *Fertil Steril* 2003;80:390–7.
13. Filicori M, Cognigni GE, Samara A, Melappioni S, Perri T, Cantelli B, et al. The use of LH activity to drive folliculogenesis: exploring uncharted territories in ovulation induction. *Hum Reprod Update* 2002;8:543–57.
14. Shoham Z, Balen A, Patel A, Jacobs HS. Results of ovulation induction using human menopausal gonadotropin or purified follicle-stimulating hormone in hypogonadotropic hypogonadism patients. *Fertil Steril* 1991;56:1048–53.
15. Couzinet B, Lestrat N, Brailly S, Forest M, Schaison G. Stimulation of ovarian follicular maturation with pure follicle-stimulating hormone in women with gonadotropin deficiency. *J Clin Endocrinol Metab* 1988;66:552–6.
16. European Recombinant Human LH Study Group. Recombinant human luteinizing hormone (LH) to support recombinant human follicle-stimulating hormone (FSH)-induced follicular development in LH- and FSH-deficient anovulatory women: a dose-finding study. *J Clin Endocrinol Metab* 1998;83:1507–14.
17. Filicori M, Cognigni GE, Taraborrelli S, Spetoli D, Ciampaglia W, de Fatis CT. Low-dose human chorionic gonadotropin therapy can improve sensitivity to exogenous follicle-stimulating hormone in patients with secondary amenorrhea. *Fertil Steril* 1999;72:1118–20.
18. Kousta E, White DM, Piazzi A, Loumaye E, Franks S. Successful induction ovulation and completed pregnancy using recombinant human luteinizing hormone and follicle stimulating hormone in a woman with Kallmann's syndrome. *Hum Reprod* 1996;11:70–1.
19. Filicori M, Cognigni GE, Pocognoli P, Ciampaglia W, Bernardi S. Current concepts and novel applications of LH activity in ovarian stimulation. *Trends Endocrinol Metab* 2003;14:267–73.
20. Nugent D, Vandekerckhove P, Hughes E, Arnot M, Lilford R. Gonadotrophin therapy for ovulation induction in subfertility associated with polycystic ovary syndrome. *Cochrane Database Syst Rev* 2000;(4):CD000410.
21. Yarali H, Bukulmez O, Gurgan F. Urinary follicle-stimulating hormone (FSH) versus recombinant FSH in clomiphene citrate-resistant, normogonadotropic, chronic anovulation: a prospective randomized study. *Fertil Steril* 1999;72:276–81.
22. Van Der Meer M, Hompes PG, De Boer JA, Schats R, Schoemaker J. Cohort size rather than follicle-stimulating hormone threshold level determines ovarian sensitivity in polycystic ovary syndrome. *J Clin Endocrinol Metab* 1998;83:423–6.
23. White DM, Polson DW, Kiddy D, Sagle P, Watson H, Gilling-Smith C, et al. Induction of ovulation with low-dose gonadotropins in polycystic ovary syndrome: an analysis of 109 pregnancies in 225 women. *J Clin Endocrinol Metab* 1996;81:3821–4.
24. Fauser BC, Van Heusden AM. Manipulation of human ovarian function: physiological concepts and clinical consequences. *Endocr Rev* 1997;18:71–106.
25. Calaf Alsina J, Ruiz Balda JA, Romeu Sarrió A, Caballero Fernández V, Cano Trigo I, Gómez Parga JL, et al. Ovulation induction with a starting dose of 50 IU of recombinant follicle stimulating hormone in WHO group II anovulatory women: the IO-50 study, a prospective, observational, multicentre, open trial. *BJOG* 2003;110:1072–7.
26. Christin-Maitre S, Hugues JN. A comparative randomized multicentric study comparing the step-up versus step-down protocol in polycystic ovary syndrome. *Hum Reprod* 2003;18:1626–31.
27. Andoh K, Mizunuma H, Liu X, Kamijo T, Yamada K, Ibuki Y. A comparative study of fixed-dose, step-down, and low-dose step-up regimens of human menopausal gonadotropin for patients with polycystic ovary syndrome. *Fertil Steril* 1998;70:840–6.
28. Balasch J, Fabregues F, Creus M, Puerto B, Penarrubia J, Vanrell JA. Follicular development and hormone concentrations following recombinant FSH administration for anovulation associated with polycystic ovarian syndrome: prospective, randomized comparison between low-dose step-up and modified step-down regimens. *Hum Reprod* 2001;16:652–6.
29. van Santbrink EJ, Fauser BC. Urinary follicle-stimulating hormone for normogonadotropic clomiphene-resistant anovulatory infertility: prospective, randomized comparison between low dose step-up and step-down dose regimens. *J Clin Endocrinol Metab* 1997;82:3597–602.
30. Chang P, Kenley S, Burns T, Denton Currie K, DeVane G, et al. Recombinant human chorionic gonadotropin (rhCG) in assisted reproductive technology: results of a clinical trial comparing two doses of rhCG (Ovidrel) to urinary hCG (Profasi) for induction of final follicular maturation in in vitro fertilization-embryo transfer. *Fertil Steril* 2001;76:67–74.
31. European Recombinant LH Study Group. Human recombinant luteinizing hormone is as effective as, but safer than, urinary human chorionic gonadotropin in inducing final follicular maturation and ovulation in in vitro fertilization procedures: results of a multicenter double-blind study. *J Clin Endocrinol Metab* 2001;86:2607–18.
32. Balasch J, Tur R, Creus M, Buxaderas R, Fabregues F, Ballesta JL, et al. Triggering of ovulation by a gonadotropin releasing hormone agonist in gonadotropin-stimulated cycles for prevention of ovarian hyperstimulation syndrome and multiple pregnancy. *Gynecol Endocrinol* 1994;8:7–12.
33. Lewit N, Kol S, Manor D, Itskovitz-Eldor J. Comparison of gonadotropin-releasing hormone analogues and human chorionic gonadotropin for the induction of ovulation and prevention of ovarian hyperstimulation syndrome: a case-control study. *Hum Reprod* 1996;11:1399–402.
34. Mulders AGMJ, Laven JSE, Eijkemans MJ, Fauser BCJM. Patient predictors for outcome of gonadotrophin ovulation induction in women with normogonadotrophic anovulatory infertility: a meta-analysis. *Hum Reprod Update* 2003;9:429–49.
35. Fluker MR, Urman B, Mackinnon M, Barrow SR, Pride SM, Ho Yuen B. Exogenous gonadotrophin therapy in World Health Organization groups I and II ovulatory disorders. *Obstet Gynecol* 1994;83:189–96.
36. Nugent D, Vandekerckhove P, Hughes E, Arnot M, Lilford R. Gonadotrophin therapy for ovulation induction in subfertility associated with polycystic ovary syndrome. *Cochrane Database Syst Rev* 2000;(3):CD000410.
37. Ron-el R, Soffer Y, Langer R, Herman A, Weintraub Z, Caspi E. Low multiple pregnancy rate in combined clomiphene citrate-human menopausal gonadotrophin treatment for ovulation induction or enhancement. *Hum Reprod* 1989;4:495–500.
38. Kemmann E, Jones JR. Sequential clomiphene citrate-menotropin therapy for induction or enhancement of ovulation. *Fertil Steril* 1983;39:772–9.
39. Legro RS, Barnhart HX, Schlaff WD, Carr BR, Diamond MP, Carson SA, et al., Cooperative Multicenter Reproductive Medicine Network. Clomiphene, metformin, or both for infertility in the polycystic ovary syndrome. *N Engl J Med* 2007;356:551–66.
40. Palomba S, Orio F Jr, Falbo A, Manguso F, Russo T, Cascella T, et al. Prospective parallel randomized, double-blind, double-dummy controlled clinical trial comparing clomiphene citrate and metformin as the first-line treatment for ovulation induction in nonobese anovulatory women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2005;90:4068–74.
41. De Leo V, la Marca A, Ditto A, Morgante G, Cianci A. Effects of metformin on gonadotropin-induced ovulation in women with polycystic ovary syndrome. *Fertil Steril* 1999;72:282–5.
42. van Santbrink EJ, Hohmann FP, Eijkemans MJ, Laven JS, Fauser BC. Does metformin modify ovarian responsiveness during exogenous FSH ovulation induction in normogonadotrophic anovulation? A placebo-controlled double-blind assessment. *Eur J Endocrinol* 2005;152:611–7.
43. Weerakiet S, Lertvikoool S, Tingthanatikul Y, Wansumrith S, Leelaphiwat S, Jultanas R. Ovarian reserve in women with polycystic ovary syndrome who underwent laparoscopic ovarian drilling. *Gynecol Endocrinol* 2007;2:1–6.

44. Farquhar C, Lilford RJ, Marjoribanks J, Vandekerckhove P. Laparoscopic "drilling" by diathermy or laser for ovulation induction in anovulatory polycystic ovary syndrome. *Cochrane Database Syst Rev* 2007;(3):CD001122.
45. Smits J, Erard P, Camus M, Devroey P, Tournaye H, Wisanto A, et al. Pituitary gonadotrophin secretory capacity during the luteal phase in superovulation using GnRH-agonists and HMG in a desensitization or flare-up protocol. *Hum Reprod* 1992;7:1225-9.
46. Beckers NG, Macklon NS, Eijkemans MJ, Ludwig M, Felberbaum RE, Diedrich K, et al. Nonsupplemented luteal phase characteristics after the administration of recombinant human chorionic gonadotropin, recombinant luteinizing hormone, or gonadotropin-releasing hormone (GnRH) agonist to induce final oocyte maturation in in vitro fertilization patients after ovarian stimulation with recombinant follicle-stimulating hormone and GnRH antagonist cotreatment. *J Clin Endocrinol Metab* 2003;88:4186-92.
47. Tavaniotou A, Devroey P. Luteal hormone profile of oocyte donors stimulated with a GnRH antagonist compared with natural cycles. *Reprod Biomed Online* 2006;13:326-30.
48. Quintero RB, Urban R, Lathi RB, Westphal LM, Dahan MH. A comparison of letrozole to gonadotropins for ovulation induction, in subjects who failed to conceive with clomiphene citrate. *Fertil Steril* 2007;88:879-85.
49. Casper RF. Aromatase inhibitors in ovarian stimulation. *J Steroid Biochem Mol Biol* 2007;106:71-5.
50. Badawy A, Metwally M, Fawzy M. Randomized controlled trial of three doses of letrozole for ovulation induction in patients with unexplained infertility. *Reprod Biomed Online* 2007;14:559-62.
51. Forman R, Gill S, Moretti M, Tulandi T, Koren G, Casper R. Fetal safety of letrozole and clomiphene citrate for ovulation induction. *J Obstet Gynaecol Can* 2007;29:668-71.
52. Tulandi T, Martin J, Al-Fadhli R, Kabli N, Forman R, Hitkari J, et al. Congenital malformations among 911 newborns conceived after infertility treatment with letrozole or clomiphene citrate. *Fertil Steril* 2006;85:1761-5.
53. Shibahara H, Kikuchi K, Hirano Y, Suzuki T, Takamizawa S, Suzuki M. Increase of multiple pregnancies caused by ovulation induction with gonadotropin in combination with metformin in infertile women with polycystic ovary syndrome. *Fertil Steril* 2007;87:1487-90.
54. Ratts VS, Pauls RN, Pinto AB, Kraja A, Williams DB, Odem RR. Risk of multiple gestation after ovulation induction in polycystic ovary syndrome. *J Reprod Med* 2007;52:896-900.
55. Stone J, Belogolovkin V, Matho A, Berkowitz RL, Moshier E, Eddleman K. Evolving trends in 2000 cases of multifetal pregnancy reduction: a single-center experience. *Am J Obstet Gynecol* 2007;197:394.
56. Practice Committee of the American Society for Reproductive Medicine. Ovarian hyperstimulation syndrome. *Fertil Steril* 2006;86(Suppl 5):S178-83.
57. Brinton LA, Moghissi KS, Scoccia B, Westhoff CL, Lamb EJ. Ovulation induction and cancer risk. *Fertil Steril* 2005;83:261-74.
58. Brinton LA, Lamb EJ, Moghissi KS, Scoccia B, Althuis MD, Mabie JE, et al. Ovarian cancer risk after the use of ovulation-stimulating drugs. *Obstet Gynecol* 2004;103:1194-203.
59. Glud E, Kjaer SK, Troisi R, Brinton LA. Fertility drugs and ovarian cancer. *Epidemiol Rev* 1998;20:237-57.